



GROW SCHEDULE



	VEGETATIVE PHASE					GENERATIVE PHASE					
	ROOTING		1		I I			П	III	III IV	
DAYS AND WEEKS	D1-5	D6-25	W1	W2	W3	W4	W5	W6	W7	W8	W9
LIGHT HOURS	18	18	12	12	12	12	12	12	12	12	12
	Dosages in ml per 10 L										
CANNA AQUA VEGA A	25	28	32	32							
CANNA AQUA VEGA B	25	28	32	32							
CANNA AQUA FLORES A					34	34	34	34	30	30	
CANNA AQUA FLORES B					34	34	34	34	30	30	
_											
CANNA RHIZOTONIC	40	20	20	20	5	5	5	5	5		
CANNA RHIZOTONIC CANNAZYM	40	20 25	20 25	20 25	5 25	5 25	5 25	5 25	5 25	25 - 50 [*]	25 - 50
4	40							-		25 - 50 * 20	25 - 50 20
CANNAZYM	40		25	25	25	25	25	25	25		
CANNAZYM CANNABOOST	40		25	25	25	25	25	25	25		
CANNAZYM CANNABOOST CANNA PK 13/14		25	25 20	25	25 20 **	25	25	25 20 15	25 20	20	20
CANNAZYM CANNABOOST CANNA PK 13/14		25	25 20	25	25 20 **	25 20 0 - 11	25	25 20 15	25 20	20	20
CANNAZYM CANNABOOST CANNA PK 13/14 CANNA CALMAG AGENT***	0 - 11	25 0 - 11	25 20 0 - 11	25 20 0 - 11	25 20 ** 0 - 11	25 20 0 - 11 mS/cm	25 20 0 - 11	25 20 15 0 - 11	25 20 0 - 11	20 0 - 11	20
CANNAZYM CANNABOOST CANNA PK 13/14 CANNA CALMAG AGENT*** EC WATER***	0 - 11	0 - 11 0.2	25 20 0 - 11	25 20 0 - 11	25 20 ** 0 - 11	25 20 0 - 11 mS/cm 0.2	25 20 0 - 11	25 20 15 0 - 11	25 20 0 - 11	0 - 11	20

This cultivation schedule is intended for plants that develop flowers or fruits, starting from a one-week-old cutting or seedling. The duration of each phase may vary depending on the species.

- * Double CANNAZYM dosage to 50 ml/10L, if substrate is reused.
- ** Increase CANNABOOST to a maximum of 40 ml/10L through flowering for extra strength
- *** CANNA CALMAG AGENT rate depends on the EC_{water}, which ideally should be 0.2-0.4 mS/cm. If the starting water has an EC of 0.0, Calmag could be applied at the maximum rate to achieve an EC of 0.4 mS/cm. If starting water is already higher than 0.0, a lower dose of Calmag should be used.

 This example schedule assumes an EC_{water} that is (adjusted with Calmag to) 0.2 mS/cm.

**** EC includes the full dose of CANNA PK 13/14 (15 ml/10 L).

For a longer growth phase, repeat the recommended dosages from Rooting period II (D6-25). For a longer flowering phase, repeat the recommended dosages from Generative Phase I (W3-W5).

Add CANNA PK 13/14 in the period 3-4 weeks before harvest.



