

# CANNA TERRA

## GROW SCHEDULE



	VEGETATIVE PHASE				GENERATIVE PHASE						
	ROOTING	I	II		I			II	III	IV	
<b>DAYS AND WEEKS</b>	D1-5	D6-25	W1	W2	W3	W4	W5	W6	W7	W8	W9
<b>LIGHT HOURS</b>	18	18	12	12	12	12	12	12	12	12	12
Dosages in ml per 10 L											
<b>CANNA TERRA VEGA</b>	41	54	62	62							
<b>CANNA TERRA FLORES</b>					66	66	66	66	66	44	
<b>CANNA RHIZOTONIC</b>	40	20	20	20	5	5	5	5	5		
<b>CANNAZYM</b>		25	25	25	25	25	25	25	25	25 - 50 *	25 - 50
<b>CANNABOOST</b>			20 **	20	20	20	20	20	20	20	20
<b>CANNA PK 13/14</b>								15			
<b>CANNA CALMAG AGENT***</b>	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11	0 - 11
mS/cm											
<b>EC WATER***</b>	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
<b>EC NUTRITION</b>	1.0	1.3	1.5	1.5	1.5	1.5	1.5	1.9 ****	1.5	1.0	
<b>EC TOTAL</b>	1.4	1.7	1.9	1.9	1.9	1.9	1.9	2.3	1.9	1.4	0.4

*This cultivation schedule is intended for plants that develop flowers or fruits, starting from a one-week-old cutting or seedling. The duration of each phase may vary depending on the species.*

- \* Double CANNAZYM dosage to 50 ml/10L, if substrate is reused.
- \*\* Increase CANNABOOST to a maximum of 40 ml/10L through flowering for extra strength.
- \*\*\* CANNA CALMAG AGENT rate depends on the EC<sub>water</sub>, which ideally should be 0.2-0.4 mS/cm. If the starting water has an EC of 0.0, Calmag could be applied at the maximum rate to achieve an EC of 0.4 mS/cm. If starting water is already higher than 0.0, a lower dose of Calmag should be used.  
This example schedule assumes an EC<sub>water</sub> that is (adjusted with Calmag to) 0.4 mS/cm.
- \*\*\*\* EC includes the full dose of CANNA PK 13/14 (15 ml/10 L).

For a longer growth phase, repeat the recommended dosages from Vegetative Phase I (D6-25). For a longer flowering phase, repeat the recommended dosages from Generative Phase I (W3-W5). Add CANNA PK 13/14 in the period 3-4 weeks before harvest.

